



**UNIVERSITI PUTRA MALAYSIA**

**SEROEPIDEMIOLOGY, DIAGNOSIS, ISOLATION AND  
CHARACTERISATION OF *NEOSPORA CANINUM* AMONG CATTLE IN  
MALAYSIA**

**CHEAH TONG SOON**

**FPV 2004 19**

**SEROEPIDEMIOLOGY, DIAGNOSIS,  
ISOLATION AND CHARACTERISATION OF  
*NEOSPORA CANINUM* AMONG CATTLE IN  
MALAYSIA**

**CHEAH TONG SOON**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALAYSIA**

**2004**



**SEROEPIDEMIOLOGY, DIAGNOSIS, ISOLATION AND  
CHARACTERISATION OF *NEOSPORA CANINUM* AMONG  
CATTLE IN MALAYSIA**

**By**

**CHEAH TONG SOON**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of  
Philosophy**

**September 2004**



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy.

**SEROEPIDEMIOLOGY, DIAGNOSIS, ISOLATION AND  
CHARACTERISATION *NEOSPORA CANINUM* AMONG CATTLE IN  
MALAYSIA**

By

**CHEAH TONG SOON**

**September 2004**

**Chairman : Associate Professor Rehana Abdullah Sani, Ph. D.**

**Faculty : Veterinary Medicine**

*Neospora caninum* is an apicomplexan protozoan parasite that has worldwide distribution and is known to cause abortion and congenital disease in cattle and dogs. This thesis deals with various aspect of investigations on neosporosis in Malaysian cattle which include seroprevalence in cattle, modes of transmission of *N. caninum* in cattle, effects of *N. caninum* infections in pregnant dairy cattle naturally infected with the parasite, isolation of *N. caninum* and comparison of the standard diagnostic technique, IFAT with ISCOM ELISA.

A cross-sectional study was conducted in dairy and beef cattle in Peninsular Malaysia and Sabah to determine the seroprevalence of *N. caninum* and the association of breeds and farm location to the seroprevalence. In Peninsular Malaysia, a total of 1531 sera from Sahiwal-Friesian on three large scale (D1, D2,

D3) and 21 small scale (D4) farms were tested for *N. caninum* antibodies using the IFAT. A total of 1413 sera from beef cattle which comprised Nelore, Droughtmaster and Kedah-Kelantan on four large scale farms (B1-B4) were also tested. In Peninsular Malaysia, the seroprevalence of *N. caninum* in the dairy cattle ranged from 1.9% in D2 farm to 6.7% in D1 farm, while in beef animals the *N. caninum* seroprevalence varied between 2.0% in B4 farm and 5.4% in B2 farm. Among the dairy cattle there was significance between the seroprevalence of *N. caninum* in D2 farm and those of D1 and D3 farms ( $p<0.05$ ). The seroprevalence of *N. caninum* in D1 farm was significantly higher than those in B1, B3 and B4 farms ( $p<0.05$ ). The *N. caninum* seroprevalence in D3 farm was significantly higher than in B4 farm. In Sabah, a total of 624 sera from Sahiwal cross and pure Friesian cattle on 6 large scale (SD1-SD6) and two small scale (SD7, SD8) farms were tested. A total of 240 sera from beef cattle which were Droughtmaster and Brahman crosses on 3 large scale (SB1-SB3) farms were tested. The seroprevalence was highest in the cattle on SD2 farm (32.5%) and lowest in the cattle on SD7 farm (2.7%). The seroprevalence of *N. caninum* of cattle on farms SB1 and SB2 were 12.5% respectively, while in SB3 farm the figure was 2.5%. There was a significant difference between the seroprevalence of *N. caninum* cattle on D2 farm and cattle on the other farms ( $p<0.05$ ).

In the investigation on modes of transmission, a cross-sectional study was conducted on the relationship between seropositivity of cows and their calves after colostral intake. Blood samples were collected from 286 dams and their respective calves on D1-D3 farms and from another 233 pairs of beef cattle on B1-B4 farms. There was a strong association between seropositivity of dams and that of their calves ( $p<0.001$ ).

The association of seroprevalence of precolostral calves with their dams was examined by collection of blood samples from 100 dams and their calves before colostral intake. Out of the 100 dams, 13 were seropositive and produced nine (69.2%) congenitally infected calves. There was strong association between presence of antibodies in the dam and congenital infection ( $p < 0.001$ ). A total of 83 seronegative heifers were also monitored prospectively from birth for evidence of post-natal infection. No evidence of seroconversion was found in the seronegative heifers.

A study conducted on D1 farm to assess the reproductive performance of eight pregnant seropositive heifers and 10 pregnant seropositive cows till pregnancy was terminated by abortion or parturition. Two heifers repeatedly returned to oestrus. Abortions were seen in four heifers. Two heifers produced weak and underweight calves and one of the calves died 12 hours after birth. The precolostral sera from the two calves had *N. caninum* antibodies. Abortion occurred in one out of the 10 pregnant cows. All the remaining nine cows produced normal calves whose precolostral and postcolostral sera were positive for *Neospora* antibodies. Paired serum samples from the aborted heifers and cow revealed no evidence of active infections due to *Br. abortus*, *Leptospira* spp., *S. dublin*, IBR virus, BVD virus or *T. gondii*. Reproductive disorders were commonly seen in the seropositive heifers when compared to seropositive cows and the difference was found to be significant ( $p < 0.05$ ).

In an attempt on isolation, brain homogenate from a congenitally infected calf born to a seropositive heifer was inoculated intraperitoneally into 10 BALB/c mice. Four out of the 10 mice died during the observation period and brains from the remaining 6 were homogenised and cultured onto Vero cell lines. Tachyzoites resembled those of *N. caninum* were detected in the cell cultures 14 days after the inoculation of brain homogenate from one of the six mice. Serology, electron microscopy and molecular studies revealed no difference between this new Malaysian bovine isolate and reference *N. caninum* isolate Nc-1, confirming that it was *N. caninum*. The isolate was designated Nc-MalB1

A total of 661 dairy and 375 beef cattle sera from Peninsular Malaysia and Sabah were used in the comparison IFAT with ISCOM ELISA in the detection of *N. caninum* antibodies. Based on the cut-offs recommended by the manufacturers, there was reasonably good agreement between the two tests as indicated by high kappas' values of 0.80 in dairy and 0.82 in beef cattle.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia,  
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah.

**SEROEPIDEMIOLOGI, DIAGNOSA, PENGASINGAN DAN  
PENCIRIAN *NEOSPORA CANINUM* PADA LEMBU DI MALAYSIA**

Oleh

**CHEAH TONG SOON**

**September 2004**

**Pengerusi : Profesor Madya Rehana Abdullah Sani, Ph.D.**

**Fakulti : Perubatan Veterinar**

*Neospora caninum* adalah parasit apicomplexan yang terdapat diseluruh dunia dan di ketahui sebagai penyebab keguguran dan penyakit kongenital pada lembu dan anjing. Tesis ini melibatkan beberapa penyiasatan berkaitan neoprosis pada lembu di Malaysia meliputi seroprevalen pada lembu, cara jangkitan penyakit, kesan jangkitan *N. caninum* pada lembu bunting, pengasingan *N. caninum* dan perbandingan IFAT dan ISCOM ELISA dalam mengesan antibodi .

Satu ‘cross-sectional study’ telah dijalankan ke atas lembu tenusu dan lembu pedaging di Semenanjung Malaysia dan Sabah untuk menentukan seroprevelensi *N. caninum* dan perkaitan di antara baka dan lokasi ladang dengan seroprevalensi. Di Semenanjung Malaysia, sejumlah 1531 serum dari lembu tenusu kacukan Sahiwal-Friesian di tiga buah ladang berskala besar (D1, D2, D3) dan 21 ladang kecil diuji



bagi mengesan kehadiran antibodi terhadap *N. caninum* menggunakan IFAT. Sejumlah 1413 serum ( $n=1413$ ) dari lembu pedaging yang terdiri dari Nelore, Droughtmaster dan Kedah-Kelantan dalam empat ladang berskala besar (B1-B4) juga diuji menggunakan kaedah yang sama. Di Semenanjung Malaysia, seroprevalensi lembu tenusu kacukan Sahiwal Friesian (SFX) adalah antara 1.9% di ladang D2 hingga 6.7% di ladang D1. Terdapat perbezaan seroprevalen yang signifikan di kalangan lembu tenusu di ladang D2 berbanding lembu di ladang D1 dan D3 ( $p<0.05$ ). Pada lembu pedaging, seroprevalen pada Droughtmaster (DM) adalah tertinggi (5.4%) dan seroprevalen pada baka indigenous Malaysia, lembu Kedah Kelantan di ladang B4 adalah terendah (2%) dan perbezaan ini adalah signifikan ( $p<0.05$ ). Diantara lembu tenusu dan lembu pedaging, seroprevalen lembu pedaging baka Nelore dan KK adalah rendah dan signifikan berbanding lembu tenusu di ladang D1 dan D3 ( $p<0.05$ ).

Di Sabah, sejumlah 624 serum dari 6 ladang tenusu berskala besar (SD1-SD6) dan 2 ladang kecil (SD7,SD8) baka SFX dan Friesian telah diuji. Seroprevalen pada ternakan di ladang SD2 (32.5%) adalah tertinggi dan terendah pada lembu di ladang SD7 (2.7%). Seroprevalen pada lembu di ladang SD2 adalah tertinggi berbanding lembu tenusu dan pedaging di Sabah dan Semenanjung Malaysia ( $p<0.001$ ). Sejumlah 240 serum lembu pedaging di 3 ladang berskala besar (SB1-SB3) telah diuji dan ternakan berbak kacukan seperti DM, Brahman (Br). Kacukan Br, Bali (B) dan kacukan B. Seroprevalen lembu di ladang SB1 dan SB2 (12.5%) adalah tinggi dan signifikan berbanding lembu di SB3 (2.5%). Seroprevalen lembu di ladang SB1 dan SB2 juga tinggi dan signifikan berbanding lembu tenusu dan lembu

pedaging di Semenanjung Malaysia ( $p<0.05$ ). Perkaitan diantara seropositif yang lebih tinggi dengan lembu SFX yang dibeli pada ladang D1 and D2 mempunyai signifikasi yang tinggi ( $p<0.001$ ).

Dalam kajian mengenal pasti cara jangkitan parasit tersebut, satu 'cross-sectional study' telah dijalankan atas perkaitan di antara status seropositif lembu betina dengan anak lembu masing-masing selepas pengambilan colostrum. Sampel darah telah diambil dari 286 lembu betina dan anak lembu dalam ladang D1-D3 serta dari 233 pasang lembu daging dalam ladang B1-B4. Terdapat perkaitan rapat di antara status seropositif lembu betina dan anak lembu ( $p<0.001$ ). Perkaitan di antara seroprevalensi anak lembu yang belum mengambil colostrum dengan lembu betina diuji melalui pengumpulan sampel darah daripada 100 lembu betina dan anak mereka sebelum pengambilan colostrum. Daripada 100 lembu betina, 13 adalah seropositif dan melahirkan 9 (69.2%) anak lembu yang mempunyai jangkitan congenital. Terdapat perkaitan yang kuat diantara kehadiran antibodi pada ibu lembu dan jangkitan kongenital ( $p<0.001$ ). Sejumlah 83 lembu dara yang seronegatif diperhatikan selepas kelahiran untuk membuktikan jangkitan postnatal. Tiada tanda – tanda serokonversi pada lembu dara yang seronegatif

Satu kajian telah dijalankan dalam ladang D1 bagi mengkaji tahap kesuburan 8 seropositif lembu dara yang mengandung pertama kali dan 10 seropositif lembu mengandung. Keguguran dilihat pada empat lembu dara bunting dan dua darinya melahirkan anak yang lemah dengan berat badan yang rendah dan seekor anak tersebut mati 12 jam selepas dilahirkan. Serum yang diambil dari dua anak lembu

sebelum mereka menghisap kolostrum mengandungi antibody terhadap *N. caninum*. Keguguran berlaku pada satu dari 10 lembu bunting. Kesemua sembilan lembu melahirkan anak lembu yang normal dengan antibodi dikesan pada darah sebelum mereka menghisap kolostrum. Serum berpasangan daripada lembu gugur menunjukkan tiada tanda jangkitan aktif disebabkan oleh *Br. Abortus*, *Leptospira* spp. *S. Dublin*, virus IBR virus, virus BVD atau *T. gondii*. Namun begitu, masalah pembiakan kerap dilihat pada lembu dara berseropositif berbanding lembu betina yang telah menghasilkan anak dan perbezaan ini adalah signifikan ( $p < 0.05$ ).

Dalam percubaan pengasingan *N. caninum*, homogenate otak daripada anak lembu berseropositif pada serum yang diambil sebelum pengambilan kolostrum, telah disuntik ke 10 BALB/c mencit secara intraperitoneal. Empat sdari 10 mencit mati dalamasa pemerhatian dan otak daripada 6 mencit yang hidup telah dihomogenasi dan dikulturkan ke dalam tisu didik vero. Tachyzoites menyerupai *N. caninum* telah dikesan di tisu didik 14 hari selepas inokulasi otak pada satu daripada enam mencit tersebut. Kajian serologi, pemerhatian elektron mikroskop dan kajian molekular menunjukkan tiada perbezaan di antara isolat baru diasingkan di Malaysia dan isolat rujukan *N. caninum* Nc-1. Isolate tersebut dikenali sebagai Nc-Ma1B1

Penyelidikan terakhir membandingkan antara IFAT dengan ISCOM ELISA bagi mengesan jangkitan *N. caninum* pada lembu. Sejumlah 661 serum lembu tenusu dan 375 lembu pedaging dari Semenanjung Malaysia dan Sabah telah digunakan bagi perbandingan. Berdasarkan 'cut –off point' vendor, terdapat persetujuan yang

munasabah di antara kedua – dua ujian tersebut yang di tunjukkan dengan nilai kappa yang tinggi iaitu 0.80 pada lembu tenusu dan 0.82 pada lembu pedaging.

## ACKNOWLEDGEMENTS

I am grateful to my supervisory committee, Assoc. Prof. Dr. Rehana Abdullah Sani (Chairperson), Dr. Chandrawathani Panchadcharam and Assoc. Prof. Dr. Mohamed Ridzwan Abdul Halim for their invaluable supervision and encouragement.

I would like to thank the Director and the staff, Institut Haiwan, Kluang, Johore, the farm Managers of all the Livestock Centres and their staff in Peninsular Malaysia and the Head of Herd Health Division and the staff, Veterinary Research Institute (VRI), Ipoh for the help in the collection of blood samples. Thanks are also extended to the Director and the staff, Livestock Disease Research Centre, Tanjung Aru, Sabah for the serum samples.

I would like to express my gratitude to the following people in VRI; Dr. Aziz Jamaluddin, the Director for his support, Dr. Sharifah Syed Hassan, the Deputy Director for facilities at the Virology Laboratory, Mr. Mohamed Zaini, Mr. Mahadi Yahaya, H. , Dr Maria, Dr Shahiruddin, Madam Sharifah Hamidah, Madam Roslina Hassan, Mr. A. Shukor Ahmad and Mr. Adnan Musbah and other staff for their assistance throughout the study.

My thanks are extended to Assoc. Prof. Dr. Fauziah Othman, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia (UPM), Serdang, Selangor,

Miss Norhayati M. M. and Mr. Ho O. K., Institute of Bioscience, UPM for the help in the transmission electron microscopy.

I am grateful to Prof. Arvid Uggla, Assoc. Prof. J. G. Mattsson and Madam E. B. Jakubek, Department of Parasitology (SWEPAR), National Veterinary Institute and Swedish University of Agricultural Sciences, Uppsala, Sweden, for the help in the molecular work and supply of cattle anti-*T. gondii* and *S. cruzi* sera under the collaborative research programme.

I wish to express my gratitude to Dr Makoto Haritani, Infectious Diseases Pathology, National Institute of Animal Health, Tsukuba, Ibaraki, Japan for his help in the immunohistochemistry.

My thanks also extended to Mr. Chua Shih Keat for his help in the statistical analyses.

I wish to express my appreciation to the Director, Institute for Medical Research, Kuala Lumpur for the supply of BALB/c mice.

The data used in this thesis was obtained while working on a project funded by the Malaysian Government through the mechanism of Intensification of Research in Priority Area Programme (01 03 02 2103) and in part by the Swedish Council and European Union Research Project QLK2-CT-2001-01050.

My special thanks go to my wife and daughter for their constant encouragement and support during the study.

I certify that an Examination Committee met on 3<sup>rd</sup> September 2004 to conduct the final examination of Cheah Tong Soon on his Doctor of Philosophy thesis entitled “Seroepidemiology, Diagnosis, Isolation and Characterisation of *Neospora caninum* among Cattle in Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Dato’ Sheikh Omar Abdul Rahman, Ph.D.**

Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

**Shaik Mohd. Amin Babjee, Ph.D.**

Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Latiffah Hassan, Ph.D.**

Lecturer  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Ooi Hong Kean, Ph.D.**

Professor  
Department of Veterinary Medicine  
National Chung Hsing University  
Taiwan  
(Independent Examiner)

---

**GULAM RUSUL RAHMAT ALI, Ph.D.**

Professor / Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

**Rehana Abdullah Sani, Ph.D.**

Associate Professor

Faculty of Veterinary Medicine

Universiti Putra Malaysia

(Chairman)

**Chandrawathani Panchadcharam, Ph.D.**

Veterinary Officer

Veterinary Research Institute

Department of Veterinary Services, Malaysia

(Member)

**Mohd. Ridzwan Abd. Halim, Ph.D.**

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

---

**AINI IDERIS, Ph.D.**

Professor / Dean

School of Graduate Studies

Universiti Putra Malaysia

Date :



## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**CHEAH TONG SOON**

Date :

# TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	vi
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF TABLES	xxi
LIST OF FIGURES	xxii
LIST OF ABBREVIATIONS	xxiv
 <b>CHAPTER</b>	
1 GENERAL INTRODUCTION	1
2 LITERATURE REVIEW	6
2.1 History and Host Distribution of <i>Neospora caninum</i>	6
2.2 Life Cycle of <i>N. caninum</i>	6
2.3 Experimental Infections	10
2.4 Neosporosis in Cattle	11
2.4.1 Clinical signs	11
2.4.2 Lesions in <i>N. caninum</i> infected foetus	13
2.4.3 Immune responses in pregnant cattle infected with <i>N. caninum</i>	14
2.4.4 Immune responses in non pregnant cattle infected with <i>N. caninum</i>	17
2.5 Prevalence and Seroepidemiology of <i>N. caninum</i> Infections in Cattle	18
2.6 Detection of Antibodies to <i>N. caninum</i>	19
2.6.1 Indirect Fluorescent Antibody Test (IFAT)	20
2.6.2 Enzyme Linked Immunosorbent Assay (ELISA)	22
2.7 Detection of <i>N. caninum</i> in Tissues	23
2.7.1 Immunohistochemical Staining of <i>N. caninum</i>	23
2.7.2 Amplication of <i>N. caninum</i> DNA	24
2.8 Isolation of <i>N. caninum</i>	24
2.9 Propagation of <i>N. caninum</i> in Mice	26
2.10 Activities of Chemotherapeutic Agents against <i>N. caninum</i> Tachyzoites	28
2.10.1 Cell Culture	28
2.10.2 In Vivo	29
2.11 Economic Impact of Neosporosis	30
2.12 Control of Neosporosis	31
2.13 Neosporosis in Horses	34

3	GENERAL MATERIALS AND METHODS	36
3.1	Farms and Management	36
3.1.1	Cattle Farms in Peninsular Malaysia	36
3.1.1.1	Dairy 1( Institute Haiwan )	37
3.1.1.2	Dairy 2 ( Air Hitam )	41
3.1.1.3	Dairy 3 ( Padang Hijau )	42
3.1.1.4	Dairy 4 ( Small scale )	42
3.1.1.5	Beef 1 ( Ulu Lepar )	43
3.1.1.6	Beef 2 ( Jelai Gemas )	43
3.1.1.7	Beef 3 ( Kuala Berang )	44
3.1.1.8	Beef 4 ( Tanah Merah )	44
3.1.2	Cattle Farms in Sabah	44
3.2	Blood Sampling	51
3.3	Indirect Fluorescent Antibody Test	51
3.3.1	Procedures	51
3.3.2	Interpretation of Results	52
3.4	ISCOM-Enzyme Linked Immunosorbent Assay	52
3.4.1	Procedures	53
3.4.2	Calculation	54
3.4.3	Interpretation of Results	54
3.5	Determination of gamma-Glutamyltransferase Activity in Serum	55
3.5.1	Procedures	55
3.5.2	Calculation	55
3.6	Latex Agglutination Test for <i>Toxoplasma gondii</i>	56
3.6.1	Procedures	56
3.6.2	Interpretation of Results	56
3.7	Data of Animals	57
3.8	Mouse Inoculation with Brain Homogenate	57
3.9	Cultivation of <i>N. caninum</i>	57
3.10	Molecular Analysis	58
3.11	Purification of a Malaysian Isolate of <i>N. caninum</i> for IFAT	59
3.12	Preparation of IFAT Slides with a Malaysian Isolate of <i>N.caninum</i>	61
3.13	Scanning Electron Microscopy ( SEM )	61
3.14	Transmission Electron Microscopy	62
3.15	Microscopic agglutination test ( MAT )	63
3.15.1	Procedures	64
3.15.2	Interpretation of Results	65
3.16	<i>Salmonella dublin</i> Serum Agglutination Test	65
3.16.1	Procedures	65
3.16.2	Interpretation of Results	66
3.17	<i>Brucella abortus</i> – Complement Fixation Test	66
3.17.1	Procedures	66
3.17.2	Interpretation of Results	67
3.18	Infectious Bovine Rhinotracheitis ( IBR ) Microtitration Neutralisation Test	68

3.18.1	Procedures	68
3.18.2	Interpretation of Results	69
3.19	Bovine Viral Diarrhoea ( BVD ) Microtitration Neutralisation Test	69
3.19.1	Procedures	70
3.19.2	Interpretation of Results	70
3.20	Histology and Immunohistochemistry	71
3.20.1	Histology	71
3.20.2	Immunohistochemistry	72
3.21	Bacteriological Examination	72
3.22	Statistical Analyses	73
4	SEROEPIDEMIOLOGY OF <i>N. CANINUM</i> INFECTION IN DAIRY AND BEEF CATTLE	74
4.1	Introduction	74
4.2	Materials and Methods	77
4.2.1	Seroprevalence of <i>N. caninum</i> Infection in Dairy and Beef Cattle in Peninsular Malaysia and Sabah	77
4.2.1.1	Study design	77
4.2.1.2	Statistical analyses	78
4.2.2	Vertical and Horizontal Transmission of <i>N. caninum</i> Infection	81
4.2.2.1	Study design	81
4.2.2.2	Determination of gamma- glutamyltransferase activity in serum	82
4.2.2.3	Statistical analyses	82
4.3	Results	83
4.3.1	Seroprevalence of <i>N. caninum</i> Infection	83
4.3.1.1	Peninsula	83
4.3.1.1a	General	83
4.3.1.1b	Dairy	83
4.3.1.1c	Beef	84
4.3.1.2	Sabah	86
4.3.1.2a	General	86
4.3.1.2b	Dairy	86
4.3.1.2c	Beef	86
4.3.2	Association of Risk Factors to the Seroprevalence of <i>N. caninum</i>	89
4.3.2.1	Association of breed and farm location to the seroprevalence of <i>N. caninum</i>	89
4.3.2.2	Association of newly imported SFX cattle from Australia on farms D1 and D2 to seroprevalence of <i>N. caninum</i>	89
4.3.3	Modes of Transmission	90
4.3.3.1	Effect of lactation number on <i>N. caninum</i> seroprevalence in pregnant and open cows	90
4.3.3.1a	General	90
4.3.3.1b	Dairy	91

	4.3.3.1c	Beef	93
	4.3.3.2	Association of seropositivity of cows and their postcolostral calves	93
	4.3.3.2a	General	93
	4.3.3.2b	Dairy	94
	4.3.3.2c	Beef	94
	4.3.3.3	Association of seropositivity of cows and their precolostral calves	96
	4.3.3.4	Investigation of seroconversion	97
	4.4	Discussion	97
5		REPRODUCTIVE PERFORMANCE IN HEIFERS AND COWS NATURALLY INFECTED WITH <i>N. CANINUM</i>	106
	5.1	Introduction	106
	5.2	Materials and Methods	107
	5.2.1	Study Design	107
	5.2.2	Statistical Analysis	109
	5.3	Results	110
	5.3.1	Heifers at D1 farm	110
	5.3.2	Heifers under Observation at VRI	112
	5.3.3	Cows at D1 Farm	114
	5.4	Discussion	115
6		ISOLATION AND CHARACTERISTION OF <i>N. CANINUM</i>	119
	6.1	Introduction	119
	6.2	Materials and Methods	120
	6.2.1	Necropsy of Calf	120
	6.2.2	Isolation	121
	6.2.3	Characterisation	121
	6.2.3.1	Amplification of isolate DNA	121
	6.2.3.2	Solid-phase sequencing	122
	6.2.3.3	Indirect fluorescent antibody test (IFAT)	122
	6.2.3.4	Transmission electron microscopy	122
	6.2.3.5	Scanning electron microscopy	122
	6.3	Results	123
	6.3.1	Calf	123
	6.3.2	Isolation	123
	6.3.3	Characterisation of Malaysian isolate	124
	6.3.3.1	Amplification of isolate DNA	124
	6.3.3.2	Solid-phase sequencing	124
	6.3.3.3	Indirect fluorescent antibody test	124
	6.3.3.4	Transmission electron microscopy	125
	6.3.3.5	Scanning electron microscopy	125
	6.4	Discussion	131
7		COMPARISON OF IFAT WITH ISCOM-ELISA IN THE DETECTION OF <i>N. CANINUM</i> ANTIBODIES IN DAIRY AND BEEF CATTLE	136

7.1	Introduction	136
7.2	Materials and Methods	137
7.2.1	Sera	137
7.2.2	Serological Testing	137
7.2.3	Calculation of Sensitivity and Specificity	138
7.2.4	Statistical Analysis	138
7.3	Results	139
7.3.1	Dairy Cattle	139
7.3.2	Beef Cattle	140
7.4	Discussion	141
8	GENERAL DISCUSSION AND CONCLUSION	144
9	PROPOSAL FOR FUTURE RESEARCH	153
	BIBLIOGRAPHY	154
	APPENDICES	174
	BIODATA OF THE AUTHOR	186
	PUBLICATIONS BASED ON STUDIES CONDUCTED IN THIS THESIS	188

## LIST OF TABLES

Table		Page
1	Successful isolations of <i>N. caninum</i> from cattle based on the country, type of samples and number of isolate	27
2	Population of cattle on dairy and beef farms in Peninsular Malaysia	38
3	Population of cattle on dairy and beef farms in Sabah	45
4	Number of animals sampled from cattle farms in Peninsular Malaysia	79
5	Number of animals sampled from each site in Sabah	80
6	Prevalence of IgG antibodies to <i>N. caninum</i> in dairy and beef cattle in Peninsular Malaysia	84
7	Prevalence of IgG antibodies to <i>N. caninum</i> in dairy and beef cattle in Sabah	87
8	Comparison of seropositivity of SFX cattle born on D1 and D2 farms with those of newly imported SFX cattle from Australia on the two farms	90
9	Cross-tabulation of <i>N. caninum</i> serological status of dams and their respective postcolostral calves on dairy and beef farms in Peninsular Malaysia	94
10	Prevalence of <i>N. caninum</i> antibodies in calves on dairy farms with respect to age at the sampling date	95
11	Prevalence of <i>N. caninum</i> antibodies in calves on beef farms in relation to age at the sampling date	95
12	Cross-classification of <i>N. caninum</i> serological status of 100 dams and their respective precolostral calves on D1 farm	96
13	Titres of <i>N. caninum</i> antibodies in aborted animals	111
14	Titres of antibodies against <i>B. abortus</i> , <i>Leptospira</i> spp., <i>S. dublin</i> , BVD and IBR virus in paired serum samples from aborted animals	111
15	Comparison of natural <i>N. caninum</i> infection in pregnant heifers and cows	115
16	Cross-classification of serological status of dairy cattle by IFAT and ISCOM-ELISA	139
17	Cross-tabulation of serological status of dairy cattle by IFAT and ISCOM-ELISA	140

## LIST OF FIGURES

Figure		Page
1	Locations of Cattle Farms in Peninsular Malaysia	46
2	Sahiwal-Friesian cross Cattle on a Large Scale Dairy Cattle Farm in Peninsular Malaysia	47
3	Sahiwal-Friesian cross Cattle on a Small Scale Dairy Cattle Farm in Peninsular Malaysia	47
4	Nelore Cattle on a Large Scale Beef Farm in Peninsular Malaysia	48
5	Droughtmaster Cattle on a Large Scale Beef Farm in Peninsular Malaysia	48
6	Kedah-Kelantan Cattle on a Large Scale Beef Farm in Peninsular Malaysia	49
7	Friesian Cattle on a Large Scale Dairy Farm in Sabah	49
8	Locations of Cattle Farms in Sabah	50
9	Prevalence of IgG Antibodies to <i>N. caninum</i> in Dairy Cattle in Peninsular Malaysia	85
10	Prevalence of IgG Antibodies to <i>N. caninum</i> in Beef Cattle in Peninsular Malaysia	85
11	Prevalence of IgG Antibodies to <i>N. caninum</i> in Dairy Cattle in Sabah	88
12	Prevalence of IgG Antibodies to <i>N. caninum</i> in Beef Cattle in Sabah	88
13	Prevalence of <i>N. caninum</i> Antibodies in Pregnant and Open Dairy Cows with Respect to their Lactation Number	92
14	Prevalence of <i>N. caninum</i> Antibodies in Pregnant and Open Beef Cows in Relation to their Lactation Number	92
15	A Pregnant Heifer Naturally Infected with <i>N. caninum</i> in an Experimental Shed	109



16	Levels of Serum Antibody in Pregnant Heifers Naturally Infected with <i>N. caninum</i>	113
17	A Calf Congenitally Infected with <i>N. caninum</i>	113
18	A Malaysian Isolate of <i>N. caninum</i> Tachyzoites	126
19	PCR Products of Isolate Nc-MalB1 Stained with Ethidium bromide in Agarose Gel	126
20	Comparison of ITS1 Sequence of Malaysian Isolate (Nc-MalB1) with that of Swedish Isolate (Nc-SweB1)	127
21	Tachyzoites of Isolate Nc-MalB1 Recognised by <i>N. caninum</i> Positive Control Serum in the IFAT	128
22	Transmission Electron-micrograph of a Tachyzoite of Isolate Nc-MalB1	128
23	Transmission Electron-micrograph of a Tachyzoite of Isolate Nc-MalB1(cross section)	129
24	Two dividing Tachyzoites of Nc-MalB1 attached to each other at One End	129
25a &b	Scanning Electron-micrograph of <i>N. caninum</i> Tachyzoites (Nc-MalB1)	130